



# Biopsy markers of airway inflammation and remodelling

P. K. JEFFERY\*, A. LAITINEN<sup>†</sup> AND P. VENGE<sup>‡</sup>

\*Imperial College School of Medicine at the Royal Brompton Hospital, London, U.K.

<sup>†</sup>Institute of Biomedicine, Department of Anatomy, University of Helsinki, Helsinki, Finland

<sup>‡</sup>Department of Medical Sciences, University Hospital, Uppsala, Sweden

Bronchial inflammation is a consistent feature of asthma and its chronicity probably determines disease progression. Clinical evaluation of drugs with potential disease-modifying activity requires measurement of their effects on the inflammatory and remodelling process using a variety of techniques including bronchial biopsy, and analysis of sputum, bronchoalveolar lavage, blood, urine and exhaled air.

Markers of the key components of the inflammatory process, such as the number and activation of T-cells, the number of mast cells, cytokine and chemokine release or gene expression, and eosinophil and neutrophil recruitment, can be determined in biopsy samples. Biopsies also allow assessment of the integrity and structure of the airway epithelium, the thickness of the reticular basement membrane and the numbers and ultrastructure of contractile cells. These and other markers may allow differentiation between subtypes of asthma patient according to atopic status and will help to distinguish asthma from chronic obstructive pulmonary disease.

Airway remodelling may be a consequence of chronic bronchial inflammation and is a characteristic of chronic asthma, particularly in severe asthma and when there is progressive decline in lung function. There are changes in the surface epithelium, reticular basement membrane, bronchial smooth muscle, blood vessels and mucous glands. Reliable markers of remodelling need to be identified to improve our ability to evaluate chronic asthma therapy.

**Key words:** asthma; inflammation; remodelling; biopsy.

RESPIR. MED. (2000) 94 (SUPPL. F), S9–S15

© 2000 HARCOURT PUBLISHERS LTD

## Introduction

Eosinophilic inflammation of the airways is common in patients with mild and moderately severe asthma and is considered to be fundamental to the functional changes that characterize the disease (1–3). Persistence of a chronic inflammatory response in asthmatic patients is thought to initiate the progressive structural change known as 'airway remodelling', which is probably a major determinant of increasing disease severity (4,5).

Consequently, pharmacotherapy which is demonstrated to have an anti-inflammatory property may prevent or slow the progress of asthma, rather than simply alleviating symptoms. As none of the clinical measures used to diagnose asthma can give precise and specific information about inflammation, more direct assessment is necessary, using techniques such as bronchial biopsy, sputum analysis, bronchoalveolar lavage (BAL) and analysis of exhaled air. By measuring changes in selected markers, for example

recruitment and function of a variety of inflammatory cell types, production of inflammatory mediators and expression of genes that control synthesis of pro-inflammatory proteins or influence the remodelling process, the underlying disease process can be monitored (6,7).

This overview considers appropriate markers for evaluation in bronchial biopsies where acute and chronic inflammation can be assessed *in situ* (8). Some of these markers may be useful in the future to evaluate the process of airway remodelling. It should be recognized, however, that interpretation of results from bronchial biopsy analysis poses challenges, not least because of the large degree of variability that exists between and within patients.

## Variation in asthmatic phenotype

One factor associated with asthma that can be measured simply is atopy. The presence of atopy is associated with inflammation, whether or not the patient presents with asthma. In one study, biopsies from atopic non-asthmatic patients had more activated (EG2<sup>+</sup>) eosinophils and a thicker reticular basement membrane (RBM) than those from healthy controls, but these changes were even more pronounced in atopic patients with asthma (9).

Received 10 July 2000 and accepted in revised form 12 July 2000. Correspondence should be addressed to: Prof. Peter K. Jeffery, Lung Pathology, Royal Brompton Hospital, Sydney Street, London, SW3 6NP, UK. Fax: +44 207 351 8435; E-mail: p.jeffery@ic.ac.uk

Eosinophilia, whilst variable, is a fundamental feature of asthma (10–13) and the associated increase in expression of the cytokines interleukin (IL)-4 and IL-13 (which promote selective vascular adhesion of these cells) appear to be of equal importance in atopic and non-atopic asthma (14). Expression of eosinophil chemokines such as eotaxin, monocyte chemoattractant protein-3 and RANTES, and of IL-4 and IL-5, the T helper-2 (Th<sub>2</sub>) cytokines, appears to be similar in the two patient sub-types (15,16); however, expression of granulocyte macrophage colony stimulating factor (GM-CSF), which has been shown to be predominantly associated with CD68<sup>+</sup> macrophages, appears to be greater in non-atopic than atopic patients (17).

In non-atopic and severe intractable asthma, neutrophil numbers appear to be significantly increased (18,19). The significance of this observation has yet to be clarified; in some patients, it may be a functional response to treatment with corticosteroids which, *in vitro*, have been shown to down-regulate apoptosis of neutrophils and thus promote their longevity (20).

The amount of the matrix glycoprotein, tenascin, in the RBM is greater in chronic and seasonal asthmatic patients than in healthy controls, and this elevation in levels appears to be more pronounced in patients with chronic disease (21). Epithelial integrity is reduced in atopic asthmatic patients compared to healthy controls and has been shown to be correlated with an increase in airway reactivity (22); significant correlations have also been observed between

epithelial integrity in atopic patients and the number of EG2<sup>+</sup> eosinophils and CD4<sup>+</sup> T-cells (18).

## Selection of markers of clinical inflammation

Many inflammatory markers can be followed in studies of asthmatic subjects; selected examples are listed in Table 1. In order to monitor inflammatory changes in atopic asthma, a marker of eosinophils is one obvious choice. EG1 and EG2, markers of eosinophilic cationic protein (ECP), are claimed to represent total and activated eosinophil counts, respectively. However, the validity of this depends on several factors including the way in which the cells are fixed and processed (23). Several studies have shown that EG2<sup>+</sup> cells may outnumber EG1<sup>+</sup> cells; weak EG2 positivity may be found in neutrophils, and macrophages may be rendered EG2-positive by ingesting apoptotic eosinophils. These antibodies can be applied to biopsy sections, or measurements of EG2<sup>+</sup> cells in fluid compartments can be obtained using flow cytometry. As eosinophils tend to cluster, their tissue distribution is not uniform, and they are frequently absent from single biopsy sections; several sections need to be assessed before a value representative of the biopsy is obtained (24,25). If biopsy is not practicable, eosinophil markers, e.g. ECF, can also be determined in blood, BAL or sputum. In biopsy,

TABLE 1. Markers of inflammation: selected examples

Tissue	Cell	Marker
Biopsies	Eosinophil	EG2, EPO
	Neutrophil	MPO, HNL
	T lymphocyte	CD3, CD4, CD8
	Activated T lymphocyte	CD25
	Macrophage	CD68
	Regulatory cytokines	IL-4, IL-5
	Chemokines	IL-8/eotaxin
	Structural changes	Thickened RBM Myofibroblast Goblet cell
Blood (serum or plasma)	Eosinophil	ECP, EPO Numbers in blood
	Neutrophil	MPO, HNL
Sputum	Cell surface markers	ICAM-1/VCAM-1
	Eosinophil	ECP, EPO
Urine	Neutrophil	HNL
	Eosinophil	EPX (EDN)
Expired air		Nitric oxide
		Hydrogen peroxide
		Carbon monoxide Ethane

ECP, eosinophil cationic protein; EPO, eosinophil peroxidase; EPX, eosinophil protein-X, also known as eosinophil-derived neurotoxin (EDN); HNL, human neutrophil lipocalin; ICAM-1, intracellular cell adhesion molecule-1; IL, interleukin; MPO, myeloperoxidase; RBM, reticular basement membrane; VCAM, vascular cell adhesion molecule-1.




FIG 1. Contribution of inter-subject variability to the total variability of CD45<sup>+</sup> cell numbers in bronchial biopsies taken from (a) asthma patients and (b) healthy individuals (unpublished data).

T-lymphocytes and their subsets and cells expressing the gene or protein for selected cytokines, e.g. IL-4, IL-5 or IL-8, can be counted, and structural changes can be measured such as epithelial integrity (22,26), thickness of the RBM and its content of tenascin, laminin and 'collagen' subtypes (21,27,28). Neutrophil markers are appropriate for non-atopic asthma, atopic asthma following allergen challenge or in severe asthma. Levels of myeloperoxidase (MPO) are twice as high in the plasma of non-atopic asthmatic patients than in atopic patients and healthy controls (unpublished data); this enzyme can also be determined in bronchoalveolar lavage fluid. In addition, human neutrophil lipocalin (HNL) is a very stable and specific neutrophil marker which can be measured in induced sputum samples or used as a neutrophil marker in tissues (29).

By selecting one key marker for each cell type, the total number of possible variables monitored can be reduced. Importantly, the variability between patients in bronchial

biopsy studies is a major contributor to total variability in the perceived levels of a given marker, as illustrated in Fig. 1 for CD45<sup>+</sup> T-cells 'unpublished data'. In studies where the variability between patients is expected to be high, reinforcement of results may be achieved by following two or more linked markers to confirm that a particular mechanistic 'axis' is affected. An example of such an axis would be the numbers of TH<sub>2</sub> cells with expression of one of the associated cytokines IL-4 or IL-5, and the numbers of eosinophils (30). Ideally, the axis chosen should be clearly linked with clinical outcomes, e.g. an improvement in airway hyper-responsiveness.

## Airway remodelling

When inflammation in the airways of asthmatic patients becomes chronic, a process of inappropriate progressive

structural changes known as airway remodelling is often present; inflammation and remodelling may be causally linked or be separate mechanistic processes (5,31). The production of matrix proteins and growth factors by populations of activated inflammatory cells, principally eosinophils, may be a first step in the cascade (32). Alternatively, a recent study suggests that airway hyper-responsiveness (AHR) may predispose to structural changes; in four patients who developed asthma 2 years after presenting with AHR but no other asthma symptoms, comparison of final and baseline biopsy assessments showed early increases in subepithelial 'fibrosis' and the number of CD25<sup>+</sup> and CD4<sup>+</sup> T cells (33). Evidence is also emerging that damage to the bronchial epithelium and a disruption of the normal repair mechanism may regulate subsequent remodelling (34). The outcomes associated with remodelling are thought to include an increase in AHR and a reduction in bronchodilator reversibility that probably results from reduced elasticity of the bronchus, increased smooth muscle mass, mucosal oedema and chronic secretion of mucus. More studies are needed to characterize the sequence of these distinct processes and the effect on them of disease-modifying drugs. Investigation of the mechanisms of various structural changes is now beginning.

### RETICULAR BASEMENT MEMBRANE (RBM)

A clearly defined change in asthmatics is thickening of the sub-epithelial RBM (22,28). The increased thickness of the RBM in asthmatic patients has been shown to be associated with increased bronchial hyper-reactivity (35,36), as well as with reduced FEV<sub>1</sub> (4) and an increased rate of exacerbations (36), but several studies have found no such correlations. Investigations of the correlation between RBM thickness and disease severity or AHR have also produced conflicting results (4,37–39). It has been suggested that activated eosinophils (EG2<sup>+</sup>) are involved in the genesis of a thickened subepithelial reticular layer, through the production of cytokines such as transforming growth factor  $\beta$  (TGF- $\beta$ ) (37), but this remains to be validated.

An additional contribution to the remodelling of extracellular matrix in chronic airway inflammation stems from imbalances between enzymes that degrade matrix proteins (matrix metalloproteinases) and endogenous inhibitors of these enzymes (tissue inhibitors of metalloproteinases; TIMPs). One example is the breakdown of elastin by neutrophil elastase; greater activity of this enzyme has been reported in asthma patients and a negative correlation has been shown with FEV<sub>1</sub> (40).

Regulation of the activity of matrix metalloproteinase MMP-9 is similarly disrupted in asthma patients; expression of the enzyme is greater than that of its tissue inhibitor and has been associated with increased RBM thickness and reduced FEV<sub>1</sub> (41).

### VASCULAR CHANGES

Increased blood flow to a site of injury and inflammation is a normal physiological response. Sustained pulmonary

hyperaemia leads to an increase in the thickness of the bronchial wall which narrows the airway lumen (42,43). One study of autopsies and resected lung samples from asthma patients showed that the density of blood vessels in the airway wall was greater than that in control samples (44). Recent biopsy studies have demonstrated that asthma patients have more vascularized bronchial tissue than controls (45,46). A study of the vascularity of the lamina propria in asthma patients receiving inhaled corticosteroids (ICS) showed no difference in the density of vessels compared with patients not on ICS, although the vessel density in all patients was correlated with both AHR and post-bronchodilator FEV<sub>1</sub> (47).

### SECRETION OF MUCUS

Hypersecretion of mucus into the airway lumen in asthmatic patients appears to be caused by both submucosal gland hypertrophy and goblet cell hyperplasia. Goblet cell hyperplasia is a prominent phenotypic change in the airway epithelium in newly diagnosed asthmatic patients (48). After 3 months of inhaled corticosteroid treatment, the epithelial structure was shown to be restored to normal ciliated epithelium, accompanied by a pronounced reduction in the number of inflammatory cells (13). One study has reported the proportion by volume of both mucin and goblet cells in the epithelium to be three times higher in asthmatic patients than in controls (49). This change may be driven by increased activity of platelet activating factor (PAF) or epidermal growth factor (EGF), both of which cause an increase in airway goblet cell number in animal models (50,51). EGF immunoreactivity in the epithelium, smooth muscle and goblet cells of autopsied bronchi from asthma patients has been shown to be greater than that from controls with no disease (52). IL-4 and IL-13 levels are also known to be raised in asthmatic patients (14) and both have been shown to stimulate goblet cell hyperplasia (53,54).

### SMOOTH MUSCLE AND THE ROLE OF GROWTH FACTORS

Increased smooth muscle mass contributes to thickening of the airway wall. This has been demonstrated by comparing autopsy samples from asthma patients and controls with no respiratory disease (44,55,56) and by quantifying the effect of smooth muscle mass on bronchoconstrictor-provoked increases in airway resistance observed in asthma patients (57).

A role for so-called 'myofibroblasts' in remodelling was first proposed when such cells, defined by immunostaining, were shown to be associated with the thickening of the RBM and deposition of fibronectin in the bronchial mucosa of asthma patients (58). The number of myofibroblasts in the mucosa of asthmatic and bronchitic patients was found to be correlated with epithelial and submucosal expression of TGF- $\beta$  produced by eosinophils and fibroblasts (59). Significant correlations have also been found between the increased number of myofibroblasts in asthma patients

compared with controls and the greater thickness of RBM containing collagen (types III and V) and tenascin (41).

Compared with controls, concentrations of TGF- $\beta$  are raised in asthmatic patients before and 24 h after allergen challenge (60). The number of 'myofibroblasts' were also shown to be markedly increased in asthmatic patients following allergen challenge (61), but these cells are likely to be distinct from those associated with RBM thickening, since they showed ultrastructural similarity to bronchial smooth muscle and are suggested to represent de-differentiated smooth muscle of the synthetic phenotype. The changes seen after allergen challenge bear similarity to the alterations described in response to vascular injury and disease. Another growth factor which can be monitored in biopsies is GM-CSF. This cytokine may play a fundamental role in the early stage of remodelling by prolonging the survival of eosinophils in the airway of asthmatic patients (62).

## Assessment of airway remodelling in clinical studies

The contribution of many processes to the phenomenon of airway remodelling in asthma patients complicates the selection of a single primary variable that would clearly signify the presence and rate of remodelling. Currently, the most favoured measurement is that of RBM thickness, but assessments of vascularity, myofibroblast numbers and phenotypic changes to airway and vascular smooth muscle by electron microscopy are becoming key research areas for the future.

Further studies on the structure of the RBM and its components in asthmatic patients and healthy controls are required to establish which components are significantly altered by the disease, and whether asthmatic patients have unique components not present in the healthy airway. Blood vessels from biopsy samples may be stained using markers of endothelium, e.g. CD31, or antibodies to type IV collagen which enable the basement membrane of the vessel wall to be visualized, so that the number of vessels per unit area and their diameter can be measured (45,46).

Other possible markers include those that measure the amounts of mucus-secreting tissue, i.e. the ratio of serous to mucous acini, gland size or goblet cell numbers in biopsy samples; analysis of mucin gene expression will also be important in the future (63,64). Once markers of remodelling have been clearly identified, the next challenge is to show that alterations in these markers are associated with changes in long-term clinical outcome.

## Conclusion

The clinical presentation of asthma and the immunopathology of airway inflammation in asthma have been well characterized. The inflammatory mechanisms responsible for the involvement of T-cells, eosinophil infiltration, TH<sub>2</sub> cytokine expression and chemokine expression by inflammatory and structural cells can be studied directly in

asthma patients by analysing bronchial biopsies. Variability of markers within the lung, airway wall and between patients, however, may reduce the reliability of estimates, and adequate sampling and careful application of quantitative technique are required. Possible distinctions between inflammation in atopic and non-atopic or in mild and severe asthma can be explored by employing specific markers such as HNL and MPO for neutrophils, and EG2 and ECP for eosinophils.

The field of chronic pulmonary inflammation is now being more intensively studied. Although the process of inappropriate airway remodelling is not yet understood, many of the contributing pathologies are now recognized and progress is beginning to be made in identifying potential markers that may be followed in long-term studies to discover more specific and effective treatments for chronic asthma in all its forms.

## References

1. Bousquet J, Chané P, Lacoste JY, *et al.* Eosinophilic inflammation in asthma. *N Engl J Med* 1990; **323**: 1033–1039.
2. Busse WW, Calhoun WF, Sedgwick JD. Mechanism of airway inflammation in asthma. *Am Rev Respir Dis* 1993; **147**: S20–S24.
3. Jeffery PK. Structural and inflammatory changes in COPD: a comparison with asthma. *Thorax* 1998; **53**: 129–136.
4. Chetta A, Foresi A, Del Denno M, Bertorelli G, Pesci A, Olivieri D. Airways remodelling is a distinctive feature of asthma and is related to severity of disease. *Chest* 1997; **111**: 852–857.
5. Bousquet J, Jeffery PK, Busse WW, Johnson M, Vignola AM. Asthma. From bronchoconstriction to airways inflammation and remodeling. *Am J Respir Crit Care Med* 2000; **161**: 1720–1745.
6. Adelroth E. How to measure airway inflammation: bronchoalveolar lavage and airway biopsies. *Can Respir J* 1998; **5** (Suppl. A): A18–A21.
7. Jeffery PK, Bousquet J. Methods for assessment of airways inflammation. *Eur Respir J* 1998; **11** (Suppl. 26): S1–S58.
8. Jeffery PK. Bronchial biopsies and airway inflammation. *Eur Respir J* 1996; **9**: 1583–1587.
9. Djukanović R, Lai CKW, Wilson JW, *et al.* Bronchial mucosal manifestations of atopy: a comparison of markers of inflammation between atopic asthmatics, atopic nonasthmatics and healthy controls. *Eur Respir J* 1992; **5**: 538–544.
10. Gleich GJ, Motojima S, Frigas E, Kephart GM, Fujisawa T, Kravis LP. The eosinophilic leukocyte and the pathology of fatal bronchial asthma: evidence for pathologic heterogeneity. *J Allergy Clin Immunol* 1987; **80**: 412–415.
11. Azzawi M, Bradley B, Jeffery PK, *et al.* Identification of activated T lymphocytes and eosinophils in bronchial biopsies in stable atopic asthma. *Am Rev Respir Dis* 1990; **142**: 1407–1413.

12. Azzawi M, Johnston PW, Majumdar S, Kay AB, Jeffery PK. T lymphocytes and activated eosinophils in asthma and cystic fibrosis. *Am Rev Respir Dis* 1992; **145**: 1477–1482.
13. Laitinen LA, Laitinen A, Haahtela T. A comparative study of the effects of an inhaled corticosteroid, budesonide, and a  $\beta_2$ -agonist, terbutaline, on airway inflammation in newly diagnosed asthma: a randomized, double-blind, parallel-group controlled trial. *J Allergy Clin Immunol* 1992; **90**: 32–42.
14. Humbert M, Durham SR, Kimmitt P, et al. Elevated expression of messenger ribonucleic acid encoding IL-13 in the bronchial mucosa of atopic and nonatopic subjects with asthma. *J Allergy Clin Immunol* 1997; **99**: 657–665.
15. Humbert M, Yung S, Corrigan C, et al. Bronchial mucosal expression of the genes encoding chemokines RANTES and MCP-3 in symptomatic atopic and nonatopic asthmatics: relationship to the eosinophil-active cytokines interleukin (IL)-5, granulocyte macrophage-colony-stimulating factor, and IL-3. *Am J Respir Cell Mol Biol* 1997; **16**: 1–8.
16. Ying S, Meng Q, Zeibecoglou K, et al. Eosinophil chemotactic chemokines (eotaxin, eotaxin-2, RANTES, monocyte chemoattractant protein-3 (MCP-3), and MCP-4), and C-C chemokine receptor 3 expression in bronchial biopsies from atopic and nonatopic (intrinsic) asthmatics. *J Immunol* 1999; **163**: 6321–6329.
17. Kotsimbos AT, Humbert M, Minshall E, et al. Upregulation of alpha GM-CSF-receptor in nonatopic asthma but not in atopic asthma. *J Allergy Clin Immunol* 1997; **99**: 666–672.
18. Amin K. The relationship between inflammation and structural changes in the airways of the lower and upper respiratory tract [dissertation]. Comprehensive summaries of Uppsala dissertations from the Faculty of Medicine 932. *Acta Universitatis Upsaliensis*, 2000.
19. Wenzel SE, Schwartz LB, Langmack EL, et al. Evidence that severe asthma can be divided pathologically into two inflammatory subtypes with distinct physiologic and clinical characteristics. *Am J Respir Crit Care Med* 1999; **160**: 1001–1008.
20. Meagher LC, Cousin JM, Seckl JR, Haslett C. Opposing effects of glucocorticoids on the rate of apoptosis in neutrophilic and eosinophilic granulocytes. *J Immunol* 1996; **156**: 4422–4428.
21. Laitinen A, Altraja A, Kämpe M, Linden M, Virtanen I, Laitinen LA. Tenascin is increased in airway basement membrane of asthmatics and decreased by an inhaled steroid. *Am J Respir Crit Care Med* 1997; **156**: 951–958.
22. Jeffery PK, Wardlaw AJ, Nelson FC, Collins JV, Kay AB. Bronchial biopsies in asthma. An ultrastructural, quantitative study and correlation with hyperreactivity. *Am Rev Respir Dis* 1989; **140**: 1745–1753.
23. Nakajima H, Loegering DA, Kita H, Kephart GM, Gleich GJ. Reactivity of monoclonal antibodies EG1 and EG2 with eosinophils and their granule proteins. *J Leukoc Biol* 1999; **66**: 447–454.
24. Sont JK, Willems LNA, Evertse CE, Hooijer R, Sterk PJ, van Krieken JHJM. Repeatability of measures of inflammatory cell number in bronchial biopsies in atopic asthma. *Eur Respir J* 1997; **10**: 2602–2608.
25. Sullivan P, Stephens D, Ansari T, Costello J, Jeffery PK. Variation in the measurements of basement membrane thickness and inflammatory cell number in bronchial biopsies. *Eur Respir J* 1998; **12**: 811–815.
26. Laitinen LA, Heino M, Laitinen A, Kava T, Haahtela T. Damage of the airway epithelium and bronchial reactivity in patients with asthma. *Am Rev Respir Dis* 1985; **131**: 599–606.
27. Altraja A, Laitinen A, Virtanen I, et al. Expression of laminins in the airways in various types of asthmatic patients: a morphometric study. *Am J Respir Cell Mol Biol* 1996; **15**: 482–488.
28. Roche WR, Beasley R, Williams JH, Holgate ST. Subepithelial fibrosis in the bronchi of asthmatic. *Lancet* 1989; **i**: 520–524.
29. Betsuyaku T, Nishimura M, Takeyabu K, et al. Netutrophil granule proteins in bronchoalveolar lavage fluid from subjects with subclinical emphysema. *Am J Respir Crit Care Med* 1999; **159**: 1985–1991.
30. Robinson DS, Hamid Q, Ying S, et al. Predominant TH2-like bronchoalveolar T-lymphocyte population in atopic asthma. *N Engl J Med* 1992; **326**: 298–304.
31. Laitinen A, Karjalainen E-M, Altraja A, Laitinen LA. Histopathologic features of early and progressive asthma. *J Allergy Clin Immunol* 2000; **105**: S509–S513.
32. Calhoun WJ, Sedgwick J, Busse WW. The role of eosinophils in the pathophysiology of asthma. *Ann N Y Acad Sci* 1991; **629**: 62–72.
33. Laprise C, Laviolette M, Boutet M, Boulet LP. Asymptomatic airway hyperresponsiveness: relationships with airway inflammation and remodelling. *Eur Respir J* 1999; **14**: 63–73.
34. Holgate ST, Lackie PM, Davies DE, Roche WR, Walls AF. The bronchial epithelium as a key regulator of airway inflammation and remodelling in asthma. *Clin Exp Allergy* 1999; **29** (Suppl. 2): 90–95.
35. Boulet LP, Laviolette M, Turcotte H, et al. Bronchial subepithelial fibrosis correlates with airway responsiveness to methacholine. *Chest* 1997; **112**: 45–52.
36. Hoshino M, Nakamura Y, Sim JJ. Expression of growth factors and remodelling of the airway wall in bronchial asthma. *Thorax* 1998; **53**: 21–27.
37. Minshall EM, Leung DY, Martin RJ, et al. Eosinophil-associated TGF-beta 1 mRNA expression and airways fibrosis in bronchial asthma. *Am J Respir Cell Mol Biol* 1997; **17**: 326–333.
38. Chu HW, Halliday JL, Martin RJ, Leung DYM, Szeffler SJ, Wenzel SE. Collagen deposition in large airways may not differentiate severe asthma from milder forms of the disease. *Am J Respir Crit Care Med* 1998; **158**: 1936–1944.
39. van Krieken JHJM, Evertse CE, Hooijer R, Willems LNA, Sterk PJ. Relationship between the inflammatory infiltrate in bronchial biopsy specimens and clinical severity of asthma in patients treated with inhaled steroids. *Thorax* 1996; **51**: 496–502.

40. Vignola AM, Bonanno A, Mirabella A, *et al.* Increased levels of elastase and alpha1-antitrypsin in sputum of asthmatic patients. *Am J Respir Crit Care Med* 1998; **157**: 505–511.
41. Hoshino M, Nakamura Y, Sim J, Shimojo J, Isogai S. Bronchial subepithelial fibrosis and expression of matrix metalloproteinase-9 in asthmatic airway inflammation. *J Allergy Clin Immunol* 1998; **102**: 783–788.
42. Mariassy AT, Gazeroglu H, Wanner A. Morphometry of the subepithelial circulation in sheep airways. *Am Rev Respir Dis* 1991; **143**: 162–166.
43. Mitzner W, Wagner E, Brown RH. Is asthma a vascular disorder? *Chest* 1995; **107**: S97–S102.
44. Kuwano K, Bosken CH, Paré PD, Bai TR, Wiggs BR, Hogg JC. Small airways dimensions in asthma and in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1993; **148**: 1220–1225.
45. Li X, Wilson JW. Increased vascularity of the bronchial mucosa in mild asthma. *Am J Respir Crit Care Med* 1997; **156**: 229–233.
46. Vrugt B, Wilson S, Bron A, Holgate ST, Djukanović R, Aalbers R. Bronchial angiogenesis in severe glucocorticoid-dependent asthma. *Eur Respir J* 2000; **15**: 1014–1021.
47. Orsida BE, Li X, Hickey B, Thien F, Wilson JW, Walters EH. Vascularity in asthmatic airways: relation to inhaled steroid dose. *Thorax* 1999; **54**: 289–295.
48. Laitinen LA, Laitinen A, Haahtela T. Airway mucosal inflammation even in patients with newly diagnosed asthma. *Am Rev Respir Dis* 1993; **147**: 697–704.
49. Ordoñez CL, Khashayar R, Ferrando RE, Wong HH, Hyde DM, Fahy JV. Increased goblet cell mucin content in mild and moderate asthma [abstract]. *Am J Respir Crit Care Med* 1998; **157**: A392.
50. Lou YP, Takeyama K, Grattan KM, *et al.* Platelet-activating factor induces goblet cell hyperplasia and mucin gene expression in airways. *Am J Respir Crit Care Med* 1998; **157**: 1927–1934.
51. Takeyama K, Dabbagh K, Lee HM, *et al.* Epidermal growth factor system regulates mucin production in airways. *Proc Natl Acad Sci USA* 1999; **96**: 3081–3086.
52. Amishima M, Munakata M, Nasuhara Y, *et al.* Expression of epidermal growth factor and epidermal growth factor receptor immunoreactivity in the asthmatic human airway. *Am J Respir Crit Care Med* 1998; **157**: 1907–1912.
53. Temann UA, Prasad B, Gallup MW, *et al.* A novel role for murine IL-4 in vivo: induction of MUC5AC gene expression and mucin hypersecretion. *Am J Respir Cell Mol Biol* 1997; **16**: 471–478.
54. Wills-Karp M, Luyimbazi J, Xu X, *et al.* Interleukin-13: central mediator of allergic asthma. *Science* 1998; **282**: 2258–2261.
55. Saetta M, Di Stefano A, Rosina C, Thiene G, Fabbri LM. Quantitative structural analysis of peripheral airways and arteries in sudden fatal asthma. *Am Rev Respir Dis* 1991; **143**: 138–143.
56. Carroll N, Elliot J, Morton A, James A. The structure of large and small airways in nonfatal and fatal asthma. *Am Rev Respir Dis* 1993; **147**: 405–410.
57. Lambert RK, Wiggs BR, Kuwano K, Hogg JC, Paré PD. Functional significance of increased airway smooth muscle in asthma and COPD. *J Appl Physiol* 1993; **74**: 2771–2781.
58. Brewster CE, Howarth PH, Djukanović R, Wilson J, Holgate ST, Roche WR. Myofibroblasts and subepithelial-fibrosis in bronchial asthma. *Am J Respir Cell Mol Biol* 1990; **3**: 507–511.
59. Vignola AM, Chanez P, Chiappara G, *et al.* Transforming growth factor-beta expression in mucosal biopsies in asthma and chronic bronchitis. *Am J Respir Crit Care Med* 1997; **156**: 591–599.
60. Redington AE, Madden J, Frew AJ, *et al.* Transforming growth factor-beta 1 in asthma. Measurement in bronchoalveolar lavage fluid. *Am J Respir Crit Care Med* 1997; **156**: 642–647.
61. Gizycki MJ, Adelroth E, Rogers AV, O'Byrne PM, Jeffery PK. Myofibroblast involvement in the allergen-induced late response in mild atopic asthma. *Am J Respir Cell Mol Biol* 1997; **16**: 664–673.
62. Park CS, Choi YS, Ki SY, *et al.* Granulocyte macrophage colony-stimulating factor is the main cytokine enhancing survival of eosinophils in asthmatic airways. *Eur Respir J* 1998; **12**: 872–878.
63. Jeffery PK, Li D. Airway mucosa: secretory cells, mucus and mucin genes. *Eur Respir J* 1997; **10**: 1655–1662.
64. Li D, Wang D, Majumdar S, *et al.* Localization and up-regulation of mucin (MUC2) gene expression in human nasal biopsies of patients with cystic fibrosis. *J Pathol* 1997; **181**: 305–310.